

*The views expressed in this editorial are those of the author(s) and do not necessarily reflect the position of the Canadian Medical Association or its subsidiaries, the journal's editorial board or the Canadian College of Neuropsychopharmacology.*

# Reconsidering brain tissue changes as a mechanistic focus for early intervention in psychiatry

Lena Palaniyappan, MD, PhD; Niron Sukumar, MD

Following a first episode of psychiatric symptoms, patients have a high risk of subsequent episodes of similar or variable nature. These subsequent recurrences characterize relapses<sup>1,2</sup> as well as comorbidity<sup>3</sup> depending on the nature of the first episode. After the first episode of psychosis, more than 50% experience a relapse within 3 years, with each relapse further increasing the risk of subsequent episodes over time.<sup>4</sup> Such recurrences are assumed to indicate a temporal progression in the underlying illness process.<sup>5</sup> The pathobiological mechanisms for these recurrences are hitherto unknown. Nevertheless, interrupting these unknown mechanistic processes has been the focus of long-term secondary prevention strategies after the first episode of serious mental illnesses such as schizophrenia,<sup>6</sup> depression<sup>7</sup> and bipolar disorder.<sup>8</sup>

Among various psychiatric disorders, secondary prevention (i.e., early intervention) strategies are best established for psychotic disorders. The rationale for the adoption of these strategies is irrefutable; early intervention in psychosis shortens the duration of personal suffering,<sup>9</sup> minimizes the psychosocial toxicity of untreated psychosis and promotes occupational achievement.<sup>10,11</sup> While the utilitarian argument is well established, the mechanistic rationale for early intervention in psychosis is built on targeting the underlying pathophysiology to arrest illness progression.<sup>12</sup> Structural brain abnormalities have been one of the most widely investigated features in this regard. Untreated illness is considered detrimental to brain structure, and early intervention is said to improve outcomes by arresting longitudinal reduction in grey matter tissue, termed “neuroprogression.”<sup>13</sup> The notion of disease modification by thwarting neuroprogression has also become the key theme for advocating early intervention in other severe mental illnesses such as bipolar disorder,<sup>14</sup> depression<sup>15</sup> and obsessive-compulsive disorder.<sup>16</sup> Despite the enthusiastic adoption of the idea of neuroprogression in the field of early intervention, several critical questions regarding this concept remain unanswered. In this editorial, we present an overview of neuroprogression in psychosis and argue for an alternative conceptual model for the neuroscience of early intervention. What follows is not a critique on the need for

early intervention, which in our opinion is indisputable for every psychiatric disorder.<sup>17</sup> Here we evaluate the mechanistic premise that is repeatedly invoked when discussing early intervention in psychiatry. While demonstrable brain changes are not required for early interventions to be effective, a fresh neuroscientific framework can accelerate treatment discovery and be impactful to deliver the next generation of transdiagnostic early intervention.<sup>18</sup>

## Is there a progressive loss of brain tissue in psychosis?

A number of longitudinal studies have been conducted now across various stages of psychosis. These studies largely concur on the occurrence of progressive loss of brain tissue in psychosis at a rate greater than expected from healthy individuals.<sup>19–21</sup> However, these longitudinal changes are spatially constrained, temporally limited to the period following the first episode, and of modest magnitude.<sup>22</sup> Furthermore, subtle progressive gain also occurs alongside progressive loss, though this gain is not concentrated to selected regions; thus, tissue increases are not picked up at the sample level, but are notable in covariance analyses.<sup>22–24</sup> This phenomenon is clearly demonstrated in a recent cross-sectional study of individual deviations from normative models of cortical thickness and white matter integrity in schizophrenia.<sup>25</sup> Whereas 79% of patients show infra-normal deviations (i.e., < 5th percentile value of age- and sex-matched healthy individuals), 46% show supra-normal deviations (i.e., > 95th percentile value) for at least one brain region. Infra-normal deviations in thickness are more common in medial prefrontal, insula and lateral temporal regions (but in only 15%–20% of patients), whereas supra-normal changes are more common in occipital and paracentral regions (affecting 3% of patients), indicating distributed changes that lack regional specificity.<sup>25</sup> Furthermore, in a recent detailed appraisal of this issue of grey matter increase using a meta-analytical network mapping approach, Mancuso and colleagues<sup>26</sup> concluded that whenever grey matter reduction occurs in one brain region, concurrent grey matter increase occurs in other brain regions

**Correspondence to:** L. Palaniyappan, 1232D, Robarts Research Institute, 1151 Richmond Street N., London, ON N6A 5B7; lpalaniy@uwo.ca

DOI: 10.1503/jpn.200172

across various psychiatric disorders. Interestingly, such grey matter changes of opposite polarity occurred in distinct functional networks. In particular, these co-alterations often involved reduced grey matter in the task-negative default-mode network (which is often observed to be “deactivated” during externally focused tasks in functional MRI studies) while concurrent increase occurred in task-positive subcortical and cognitive control regions.<sup>26</sup> Taken together, these findings indicate that the phenomenon commonly considered to be neuroprogression may instead be a well-coordinated compensatory process operating across various psychiatric disorders.

### Does neuroprogression indicate an unfavourable outcome?

The presence of lower grey matter volume at the outset of the first episode, even before treatment initiation, has been shown to be predictive of poor subsequent response.<sup>27–29</sup> Additionally, patients who have a longer duration of active symptoms show more longitudinal volume loss than those who are in remission.<sup>30–32</sup> Despite these reports, a causal relationship between postonset tissue loss and subsequent cognitive, functional or symptomatic decompensation is weakened by several observations. First, there is a notable lack of diagnostic specificity for progressive tissue loss. Relapses of various nature are associated with greater tissue loss, as reported in recurrent depressive disorder,<sup>33</sup> alcohol use disorder,<sup>34</sup> bipolar disorder,<sup>35</sup> migraine,<sup>36</sup> epilepsy<sup>37</sup> and chronic pain.<sup>38</sup> While there are likely differences in the affected brain regions among these illnesses, progressive tissue loss in these disorders is not suggestive of a decompensating trajectory. Second, the association between illness severity and grey matter loss is inconsistent. Several studies indicate a lack of association between tissue reduction and illness severity (as reviewed previously<sup>22,39</sup>). Some studies even indicate a reverse relationship, associating unfavourable outcomes to an increase in grey matter as opposed to a decrease.<sup>40–42</sup> Third, patients who are more symptomatic tend to receive higher doses of psychotropic medications, specifically antipsychotics, which hasten grey matter reduction.<sup>43–45</sup> Observational studies cannot resolve this confound. Hence, we cannot confidently conclude that neuroprogression is indicative of an unfavourable outcome in psychosis.

### Does treatment delay worsen neuroprogression?

Many studies that report progressive tissue loss include participants who receive clinical care at specially constituted early intervention services, indicating that neuroprogression occurs despite early psychosocial and medical care.<sup>46</sup> While experimental reduction of the duration of untreated psychosis (DUP) improves functional outcomes,<sup>47</sup> observational studies relating duration of untreated illness to tissue loss are inconclusive.<sup>19,21,43,48,49</sup> A systematic review by Anderson and colleagues<sup>50</sup> examined 43 studies that investigated the association between duration of untreated psychosis and brain structure (imaged with a variety of modalities). Of these, only 8 studies showed statistically significant findings and, even

among the brain regions identified as having significant structural associations, these findings were inconsistent across studies. Observational studies linking DUP to brain changes are limited because DUP is a complex variable affected by service availability as well as personal and family profiles. In addition, multiple measurement confounds have been neglected in prior studies. In particular, MRI studies have not separated cortical thickness, a malleable morphometric feature that is sensitive to plastic changes in adult life, from surface area or gyrification, which are largely determined during early brain development and show much smaller magnitude of changes in adult life.<sup>51</sup> A recent multimodal imaging study in medication-naïve first-episode psychosis reported that longer DUP relates to both reduced surface area and increased cortical thickness.<sup>52</sup> This contradicts the neuroprogressive hypothesis, which would predict larger reductions in brain features that are more malleable in adult life, such as cortical thickness, as opposed to features that are more developmentally determined. Rather than treatment delay causing structural changes, these findings are better explained by the fact that patients with developmentally determined structural deficits are more likely to receive delayed treatment, likely because of the insidious onset and the lack of a dramatic drop in functional levels given their premorbid deficits. This idea is supported by a large body of literature linking the indicators of developmental aberrations such as familial loading of illness risk,<sup>53</sup> poor premorbid adjustment,<sup>54–56</sup> early age of onset<sup>56</sup> and neurologic soft signs<sup>57</sup> with longer treatment delay. A prospective assessment of the same cohort, subjected to a 16-week trial of risperidone, revealed longer DUP to predict poorer treatment response.<sup>52</sup> While treatment delay diminishes the probability of favourable response, the relationship between these variables is unlikely to be mediated by progressive grey matter reduction. It is likely that the treatments we provide do not work well for latecomers,<sup>17</sup> who are likely to be those with developmental deficits.<sup>58,59</sup>

### Neuroprogression: Illness or treatment effect?

A series of observational studies have implicated antipsychotics, especially in higher doses, in neuroprogression.<sup>39,44,60</sup> However, in clinical practice, patients who receive higher doses of antipsychotics are also often those with higher symptom burden and longer illness duration. Untangling the treatment from illness effects have been hitherto impossible, but 3 recent studies provide some clarity.

Liu and colleagues<sup>61</sup> recruited a rare sample of patients with nearly 20 years of untreated schizophrenia from rural Western China. When 2 matched groups of treated patients (risperidone or clozapine) with a similar duration of illness were compared, pronounced grey matter reduction was seen in the cohort treated with antipsychotics compared with the treatment-naïve cohort. The treatment-naïve sample did exhibit widespread grey matter reduction, but also grey matter increases in specific brain regions compared with age-matched healthy controls. Being a case-control study, the temporal association between treatment administration and tissue loss cannot be inferred.

In the Staged Treatment and Acceptability Guidelines in Early Psychosis Study (STAGES), Chopra and colleagues<sup>62</sup> randomized 62 patients to antipsychotic treatment or placebo, while providing psychosocial interventions to both arms. Three months after treatment, the medicated group had an increase in pallidal volume (likely related to D2-blockade effect), but the placebo group had a reduction in pallidal volume with a small effect size difference between the 2 arms. Interestingly, at 12 months, there were no differences between the 2 groups. Given the ethical constraints of a placebo-controlled study, this sample included only patients with short DUP (< 6 mo), only a low dose of antipsychotic was used, and patients showing poor response were removed from the triple-blind experiment, thus affecting generalizability.

In the Study of the Pharmacotherapy of Psychotic Depression II (STOP-PD II), Voineskos and colleagues<sup>63</sup> randomized 88 patients with psychotic depression who achieved 8-week remission from psychosis on sertraline and olanzapine, to either discontinue olanzapine (placebo arm) or continue for 36 weeks. The olanzapine group experienced a significant reduction in cortical thickness, but not surface area. In the placebo arm, relapses were also associated with thickness reduction, albeit with smaller effect sizes than olanzapine exposure. Interestingly, in the olanzapine arm, reduction in thickness was more likely in those who sustained remission than in those who relapsed. In the olanzapine arm, reductions were predominantly in the lateral frontotemporal cortex (pars opercularis and middle temporal region), but regional differences in relapse- compared with remission-related thickness reduction were not examined. Patients in this study had psychotic depression and an average age of 55 years. The generalizability of this observation to patients with other forms of psychosis, who tend to be much younger at presentation, is unclear.

From these reports, we can conclude that a notable portion of MRI-based grey matter reduction is antipsychotic related, while illness-related reduction also occurs and relates to the periods of active symptoms (relapses). Taken together, these observations challenge the neuroprogressive interpretation. Grey matter reduction occurs in the placebo arm in both randomized controlled trials (RCTs; though only for the first 3 mo in the data reported by Chopra and colleagues<sup>62</sup>), consistent with neuroprogression. But grey matter reduction is also associated with the administration of demonstrably efficacious treatment in both RCTs as well as in the chronic untreated sample reported by Liu and colleagues<sup>61</sup>; this would not be expected if it is the result of a progressive pathophysiology. Given the larger effect size seen in the treatment arm of STOP-PD II, one may posit that treatment itself somehow hastens neuroprogression despite demonstrating superior outcomes. However, this conclusion is contradicted by the fact that in the olanzapine arm, reduction in thickness was more likely in those who experienced sustained remission than in those who relapsed. This is the reverse of what was seen in the placebo arm, where thickness reduction was shown to be more likely in patients who relapsed. Furthermore, in the STAGES study, 12 months after treatment initiation, there were no notable differences between illness-related and treatment-related changes. While it is possible

that the illness and its treatment affect the structure of different brain regions, all reductions in cortical thickness, irrespective of their regional localization, are considered deleterious under the premise of neuroprogression. If we accept the neuroprogressive interpretation, we cannot reconcile the contradiction that treatment-related structural changes appear protective, while the illness-related changes appear to be deleterious (as observed in STOP-PD II), with both processes resulting in similar brain anatomy over a longer period of time (as observed in STAGES).

How can we then explain the observation that both the illness and its treatment produce a similar change in the presumed pathophysiology? We propose that the observed progressive structural changes are reflective of physiologic adaptation rather than primary pathophysiology, and these compensatory efforts are facilitated, not alleviated, by treatment.

## Neuroprogression as an adaptive change

When we start considering neuroprogression as an adaptive change, a number of paradoxical observations in the field start to fall in place. First, when considering long-term trajectories, progressive tissue loss does not reflect longitudinal worsening; in fact, patients treated with antipsychotics get better despite tissue loss.<sup>64,65</sup> Clozapine, arguably the most effective intervention for schizophrenia, increases the progressive tissue loss despite clinical and functional improvement.<sup>66-68</sup> Second, progressive tissue loss is not specific to psychosis or to any psychiatric syndrome or individual symptom. A large amount of the variance in structural changes is explained by a single *p* factor that captures the overall burden of psychopathology.<sup>69</sup> Many brain regions are altered by the majority of brain diseases, albeit to a variable extent.<sup>70,71</sup> Third, most patients with psychosis display a pattern of cortical thickness that is indistinguishable from that of healthy controls,<sup>25,72,73</sup> indicating that neuroprogression is an unlikely mechanistic process operating in a large majority of functionally disabled patients who need active interventions. If this is the case, reversing or arresting these brain changes may not bring the desired effects of early intervention.

The extant literature reviewed here indicates that putative neuroprogression is unlikely the primary pathophysiological process to be targeted for secondary prevention efforts in psychiatry. If we refrain from invoking the framework of a progressive pathophysiology, then the mechanistic basis for the onset (i.e., first episode) and later recurrences could be considered separable. In other words, the inefficient adaptive pathobiological processes that lead to recurrences (relapses or comorbidity) are likely distinguishable from the mechanisms triggering the onset.

## Neuroadaptation as the mechanistic focus of early intervention

In many physiologic conditions, grey matter reduction confers an adaptive advantage (e.g., developmental cortical thinning<sup>74,75</sup> or pregnancy<sup>76</sup>). A higher rate of change of cortical thickness is associated with greater intelligence in healthy adolescents,<sup>77</sup> while pregnancy-associated grey matter reduction is

associated with improved maternal bonding.<sup>76</sup> While such an assertion in pathological states may be iconoclastic, it is important to recognize both benefits and costs of grey matter changes in psychiatric disorders before intervening to “protect” against them. Preclinical studies indicate that morphological changes induced by developmental adversity may indeed prime the organism for a more strategic response in the face of further adversity.<sup>78</sup> In this context, the term “neuroprogression” for post-onset structural changes in psychiatric disorders stands out as an unfortunate misnomer. A more accurate description of these MRI-based grey matter changes would be “cortical reorganization” or “structural neuroadaptation.”<sup>79</sup>

Physiologic adaptation to perturbation is a feature of all complex biological systems, including the human brain.<sup>80</sup> While restoration of pre-perturbation parameters (healthy baseline) in patients is naturally the most desirable state, a chaotic response resulting in alternate states of stability is more likely with intense, protracted perturbations<sup>81,82</sup> (as in a psychotic episode). Such “exploratory” neuroadaptation may result in distributed, albeit small effect size changes in the brain structure and may continue until a new “steady state” is reached, which may be less than optimal<sup>82</sup> (inefficient compensation or maladaptive steady state<sup>83</sup>). In the context of systems biology, psychiatric phenomena can be seen as emergent properties;<sup>84,85</sup> their appearance as well as resolution cannot be fully solved by focusing on the properties of single constituent markers, such as MRI-based brain structural changes. A deeper understanding of this exploratory adaptive response is a requisite step for devising empirically informed secondary prevention strategies. In this case, intervention to modify illness trajectories would be different from the initial treatment and would focus on guiding more adaptive biological processes to emerge in an individual.

## Investigating neuroadaptation in psychosis

The conventional approach in the field tends to view structural changes as a pathological outcome to be targeted for intervention (i.e., neuroprogression). In contrast, the systems perspective<sup>82</sup> acknowledges the inherently dynamic nature of brain structure, its interconnectedness with illness-related factors at various levels and the variations that may result from internal regulation to reduce the impact of external perturbations.<sup>86</sup> In this view, MRI-based structural changes are likely nested within a system of physiologic changes occurring at the molecular, cellular, tissue or organ (i.e., whole brain) level as well as the bodily changes that occur elsewhere in the course of psychosis (e.g., cardiometabolic changes). In addition to this interconnected system of biological changes, causal interactions also occur outside the body, at the level of individual and collective behaviour,<sup>87</sup> affecting the brain structure. As with other systems concepts, investigations of neuroadaptation require multi-scale and multi-level measures that employ network theory. At the level of brain structure, investigations that parse the relationship between various spatial units and tissue types are needed (e.g., graphs of concomitant grey matter and white matter changes, relationship between grey matter reduction across different brain areas). These structural graphs depicting an observed state of interaction must then be studied in con-

junction with graphs capturing the emergent properties of interacting signals of brain function (e.g., co-occurring changes in functional activity and symptom-based, cognitive and behavioural networks<sup>88</sup>). Crucially, as neuroadaptation results from autoregulatory, self-organizing processes, they are likely not defined by discrete, observable events but occur gradually over time. Characterizing dynamic neuroadaptation processes will require multi-level, multimodal measurements to be repeatedly captured over time.

An implicit speculation that arises from the proposal here is that antipsychotics may assist the process of neuroadaptation. The evidence to date, in our view, supports the notion that post-onset thickness change (reduction and increase) may indeed be an adaptive process. There is also emerging evidence that antipsychotics contribute to thickness reduction. Nevertheless, there is no evidence to assume that the antipsychotic-induced thickness reduction per se is therapeutically beneficial. In due course, the antipsychotic- versus illness-related thickness changes may turn out to be distinct in their spatial distribution across the brain, the underlying histological changes (e.g., increased intracortical myelination v. grey matter atrophy), the cellular constituents affected (e.g., glial v. neuronal changes), the time course (continuous or time-limited effects) as well as the potential reversibility. None of this information is presently known. Experimental “on-off” studies of antipsychotics in selected patients with multiple within-subject observations and careful characterization of symptom and functional changes are required to parse these phenomena further. Noninvasive experiments that induce short-term adaptive changes (e.g., sensory- or motor-deprivation paradigms<sup>89</sup>) may be feasible in both medicated and unmedicated patients. Such experiments will provide clarity on whether antipsychotics assist or interfere with presumed adaptive processes. Until such evidence is available, antipsychotic-induced structural changes should be considered with caution.

There is a clear need for energetically promoting early intervention in psychiatry. The psychosocial toxicity of untreated psychiatric illness is irrefutable; this forms the sufficient premise to vigorously promote early intervention. Treatment delay in psychiatric disorders occurs due to systemic factors such as chronic underinvestment in effective service delivery, societal factors pertaining to stigma, and personal health belief models. Addressing these issues remains the primary and urgent goal of secondary prevention in psychiatry. Early intervention provides substantial benefits at a program level,<sup>50</sup> but our arsenal for providing effective, minimally invasive and acceptable early intervention for the patients with varying illness severity that we see at our clinics remains inadequate. Linking early intervention to deleterious neuroprogression is untenable at best, and may hamper the progress of further enquiry into mechanistically informed secondary prevention in psychiatry. It is time for the neuroscience of early intervention to take a decisive turn away from the dark alley of neuroprogression.

**Affiliations:** From the Robarts Research Institute, Western University (Palaniyappan); the Department of Psychiatry, Western University (Palaniyappan, Sukumar); the Lawson Health Research Institute, Imaging Division (Palaniyappan); and the Department of Medical Biophysics, Western University (Palaniyappan), London, Ont., Canada.



**Acknowledgement:** L. Palaniyappan acknowledges support from the Tanna Schulich Endowed Chair of Neuroscience and Mental Health.

**Competing interests:** L. Palaniyappan reports personal fees from Janssen Canada, Otsuka Canada, SPMM Course Limited, UK, and the Canadian Psychiatric Association; book royalties from Oxford University Press; investigator-initiated educational grants from Sunovion, Janssen Canada, and Otsuka Canada; and travel support from Boehringer Ingelheim and Magstim Limited, outside the submitted work. He is a member of the *JPN* editorial board, but was not involved in the review or the decision to accept this manuscript for publication. No other competing interests were declared.

## References

- Emsley R, Chiliza B, Asmal L, et al. The nature of relapse in schizophrenia. *BMC Psychiatry* 2013;13:50.
- Andrews G. Should depression be managed as a chronic disease? *BMJ* 2001;322:419-21.
- Plana-Ripoll O, Pedersen CB, Holtz Y, et al. Exploring comorbidity within mental disorders among a Danish national population. *JAMA Psychiatry* 2019;76:259-70.
- Alvarez-Jimenez M, Priede A, Hetrick SE, et al. Risk factors for relapse following treatment for first episode psychosis: a systematic review and meta-analysis of longitudinal studies. *Schizophr Res* 2012;139:116-28.
- McGorry PD, Killackey E, Yung A. Early intervention in psychosis: concepts, evidence and future directions. *World Psychiatry* 2008;7:148-56.
- Zipursky RB, Agid O, Remington G. Improving outcomes in schizophrenia by preventing early relapses. *The Lancet Psychiatry* 2018;5:384-6.
- Nemeroff CB. The state of our understanding of the pathophysiology and optimal treatment of depression: Glass half full or half empty? *Am J Psychiatry* 2020;177:671-85.
- Post RM. Preventing the malignant transformation of bipolar disorder. *JAMA* 2018;319:1197-8.
- Albert N, Weibell MA. The outcome of early intervention in first episode psychosis. *Int Rev Psychiatry* 2019;31:413-24.
- Windell D, Norman R, Malla AK. The personal meaning of recovery among individuals treated for a first episode of psychosis. *Psychiatr Serv* 2012;63:548-53.
- Norman RMG, Mallal AK, Manchanda R, et al. Does treatment delay predict occupational functioning in first-episode psychosis? *Schizophr Res* 2007;91:259-62.
- Lieberman JA, Small SA, Girgis RR. Early detection and preventive intervention in schizophrenia: from fantasy to reality. *Am J Psychiatry* 2019;176:794-810.
- Kapczynski F, Streib LG, Kapczynski F, et al. Neuroprogression and staging in psychiatry: historical considerations. *Braz J Psychiatry* 2014;36:187-8.
- Keramatian K, Su W, Saraf G, et al. Preservation of gray matter volume in early stage of bipolar disorder: a case for early intervention [Epub ahead of print]. *Can J Psychiatry* 2020 May 18;0706743720927827. doi: 10.1177/0706743720927827.
- Moylan S, Maes M, Wray NR, et al. The neuroprogressive nature of major depressive disorder: pathways to disease evolution and resistance, and therapeutic implications. *Mol Psychiatry* 2013;18:595-606.
- Fineberg NA, Dell'Osso B, Albert U, et al. Early intervention for obsessive compulsive disorder: an expert consensus statement. *Eur Neuropsychopharmacol* 2019;29:549-65.
- Palaniyappan L, Krishnadas R. Treatment delay in early psychosis: not a linear problem. *Lancet Psychiatry* 2020;7:563-5.
- McGorry PD, Mei C. Early intervention in youth mental health: progress and future directions. *Evid Based Ment Health* 2018;21:182-4.
- Vita A, Peri LD, Deste G, et al. Progressive loss of cortical gray matter in schizophrenia: a meta-analysis and meta-regression of longitudinal MRI studies. *Transl Psychiatry* 2012;2:e190.
- Olabi B, Ellison-Wright I, McIntosh AM, et al. Are there progressive brain changes in schizophrenia? A meta-analysis of structural magnetic resonance imaging studies. *Biol Psychiatry* 2011;70:88-96.
- Fusar-Poli P, Smieskova R, Kempton MJ, et al. Progressive brain changes in schizophrenia related to antipsychotic treatment? A meta-analysis of longitudinal MRI studies. *Neurosci Biobehav Rev* 2013;37:1680-91.
- Palaniyappan L. Progressive cortical reorganisation: A framework for investigating structural changes in schizophrenia. *Neurosci Biobehav Rev* 2017;79:1-13.
- Palaniyappan L, Das T, Dempster K. The neurobiology of transition to psychosis: clearing the cache. *J Psychiatry Neurosci* 2017;42:294-9.
- Palaniyappan L, Das TK, Winmill L, et al. Progressive post-onset reorganisation of MRI-derived cortical thickness in adolescents with schizophrenia. *Schizophr Res* 2019;208:477-8.
- Ly J, Di Biase M, Cash RHF, et al. Individual deviations from normative models of brain structure in a large cross-sectional schizophrenia cohort [Epub ahead of print]. *Mol Psychiatry* 2020 Sept. 22. doi: 10.1038/s41380-020-00882-5.
- Mancuso L, Fornito A, Costa T, et al. A meta-analytic approach to mapping co-occurrent grey matter volume increases and decreases in psychiatric disorders. *Neuroimage* 2020;222:117220.
- Takahashi T, Kido M, Sasabayashi D, et al. Gray matter changes in the insular cortex during the course of the schizophrenia spectrum. *Front Psychiatry* 2020;11:659.
- Li M, Li X, Das TK, et al. Prognostic utility of multivariate morphometry in schizophrenia. *Front Psychiatry* 2019;10:245.
- Zhang X, Zhang Y, Liao J, et al. Progressive grey matter volume changes in patients with schizophrenia over 6 weeks of antipsychotic treatment and their relationship to clinical improvement. *Neurosci Bull* 2018;34:816-26.
- Rosa PGP, Zanetti MV, Duran FLS, et al. What determines continuing grey matter changes in first-episode schizophrenia and affective psychosis? *Psychol Med* 2015;45:817-28.
- Lepage M, Makowski C, Bodnar M, et al. Do unremitted psychotic symptoms have an effect on the brain? A two year follow up imaging study in first episode psychosis. *Schizophr Bull Open* 2020;1:sgaa039.
- Andreasen NC, Liu D, Ziebell S, et al. Relapse duration, treatment intensity, and brain tissue loss in schizophrenia: a prospective longitudinal MRI study. *Am J Psychiatry* 2013;170:609-15.
- Zaremba D, Dohm K, Redlich R, et al. Association of brain cortical changes with relapse in patients with major depressive disorder. *JAMA Psychiatry* 2018;75:484-92.
- Beck A, Wüstenberg T, Genauck A, et al. Effect of brain structure, brain function, and brain connectivity on relapse in alcohol-dependent patients. *Arch Gen Psychiatry* 2012;69:842-52.
- Kozicky J-M, McGirr A, Bond DJ, et al. Neuroprogression and episode recurrence in bipolar I disorder: a study of gray matter volume changes in first-episode mania and association with clinical outcome. *Bipolar Disord* 2016;18:511-9.
- Messina R, Rocca MA, Colombo B, et al. Gray matter volume modifications in migraine: a cross-sectional and longitudinal study. *Neurology* 2018;91:e280-92.
- Alvim MKM, Coan AC, Campos BM, et al. Progression of gray matter atrophy in seizure-free patients with temporal lobe epilepsy. *Epilepsia* 2016;57:621-9.
- Rodriguez-Raecke R, Niemeier A, Ihle K, et al. Structural brain changes in chronic pain reflect probably neither damage nor atrophy. *PLoS One* 2013;8:e54475.
- Emsley R, Asmal L, du Plessis S, et al. Brain volume changes over the first year of treatment in schizophrenia: relationships to antipsychotic treatment. *Psychol Med* 2017;47:2187-96.
- DeLisi LE, Stritzke P, Riordan H, et al. The timing of brain morphological changes in schizophrenia and their relationship to clinical outcome. *Biol Psychiatry* 1992;31:241-54.
- Gur RE, Turetsky BJ, Cowell PE, et al. Temporolimbic volume reductions in schizophrenia. *Arch Gen Psychiatry* 2000;57:769-75.
- Dukart J, Smieskova R, Harrisberger F, et al. Age-related brain structural alterations as an intermediate phenotype of psychosis. *J Psychiatry Neurosci* 2017;42:307-19.
- Ho B-C, Andreasen NC, Ziebell S, et al. Long-term antipsychotic treatment and brain volumes: a longitudinal study of first-episode schizophrenia. *Arch Gen Psychiatry* 2011;68:128-37.
- Guo JY, Huhtaniska S, Miettinen J, et al. Longitudinal regional brain volume loss in schizophrenia: relationship to antipsychotic medication and change in social function. *Schizophr Res* 2015;168:297-304.
- Dorph-Petersen K-A, Pierri JN, Perel JM, et al. The influence of chronic exposure to antipsychotic medications on brain size before and after tissue fixation: a comparison of haloperidol and olanzapine in macaque monkeys. *Neuropsychopharmacology* 2005;30:1649-61.

46. Théberge J, Williamson KE, Aoyama N, et al. Longitudinal grey-matter and glutamatergic losses in first-episode schizophrenia. *Br J Psychiatry* 2007;191:325-34.
47. Hegelstad WTV, Larsen TK, Auestad B, et al. Long-term follow-up of the trends pharmacol sci early detection in psychosis study: effects on 10-year outcome. *Am J Psychiatry* 2012;169:374-80.
48. Rapp C, Canela C, Studerus E, et al. Duration of untreated psychosis/illness and brain volume changes in early psychosis. *Psychiatry Res* 2017;255:332-7.
49. Goff DC, Zeng B, Ardekani BA, et al. Association of hippocampal atrophy with duration of untreated psychosis and molecular biomarkers during initial antipsychotic treatment of first-episode psychosis. *JAMA Psychiatry* 2018;75:370-378.
50. Anderson KK, Norman R, MacDougall A, et al. Effectiveness of early psychosis intervention: comparison of service users and non-users in population-based health administrative data. *Am J Psychiatry* 2018;175:443-52.
51. Storsve AB, Fjell AM, Tamnes CK, et al. Differential longitudinal changes in cortical thickness, surface area and volume across the adult life span: regions of accelerating and decelerating change. *J Neurosci* 2014;34:8488-98.
52. Maximo JO, Nelson EA, Armstrong WP, et al. Duration of untreated psychosis correlates with brain connectivity and morphology in medication-naïve patients with first-episode psychosis. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2020;5:231-8.
53. Esterberg M, Compton M. Family history of psychosis negatively impacts age at onset, negative symptoms, and duration of untreated illness and psychosis in first-episode psychosis patients. *Psychiatry Res* 2012;197:23-8.
54. Maccabe JH, Aldouri E, Fahy TA, et al. Do schizophrenic patients who managed to get to university have a non-developmental form of illness? *Psychol Med* 2002;32:535-44.
55. Norman RMG, Malla AK, Manchanda R. Early premorbid adjustment as a moderator of the impact of duration of untreated psychosis. *Schizophr Res* 2007;95:111-4.
56. Larsen TK, Friis S, Haahr U, et al. Premorbid adjustment in first-episode non-affective psychosis: distinct patterns of pre-onset course. *Br J Psychiatry* 2004;185:108-15.
57. Chen EY-H, Hui CL-M, Chan RC-K, et al. A 3-year prospective study of neurological soft signs in first-episode schizophrenia. *Schizophr Res* 2005;75:45-54.
58. Palaniyappan L, Marques TR, Taylor H, et al. Cortical folding defects as markers of poor treatment response in first-episode psychosis. *JAMA Psychiatry* 2013;70:1031-40.
59. Palaniyappan L, Marques TR, Taylor H, et al. Globally efficient brain organization and treatment response in psychosis: a connectomic study of gyrification. *Schizophr Bull* 2016;42:1446-56.
60. Ho B-C, Andreasen NC, Nopoulos P, et al. Progressive structural brain abnormalities and their relationship to clinical outcome: a longitudinal magnetic resonance imaging study early in schizophrenia. *Arch Gen Psychiatry* 2003;60:585-94.
61. Liu N, Xiao Y, Zhang W, et al. Characteristics of gray matter alterations in never-treated and treated chronic schizophrenia patients. *Transl Psychiatry* 2020;10:136.
62. Chopra S, Fornito A, Francy S, et al. Differentiating the effect of medication and illness on brain volume reductions in first-episode psychosis: a longitudinal, randomized, triple-blind, placebo-controlled MRI study. *medRxiv* 2020;2020.03.18.20038471.
63. Voineskos AN, Mulsant BH, Dickie EW, et al. Effects of antipsychotic medication on brain structure in patients with major depressive disorder and psychotic features: neuroimaging findings in the context of a randomized placebo-controlled clinical trial. *JAMA Psychiatry* 2020;77:674-83.
64. Zipursky RB, Reilly TJ, Murray RM. The myth of schizophrenia as a progressive brain disease. *Schizophr Bull* 2013;39:1363-72.
65. Lawrie SM. Are structural brain changes in schizophrenia related to antipsychotic medication? A narrative review of the evidence from a clinical perspective. *Ther Adv Psychopharmacol* 2018;8:319-26.
66. Mattai A, Chavez A, Greenstein D, et al. Effects of clozapine and olanzapine on cortical thickness in childhood-onset schizophrenia. *Schizophr Res* 2010;116:44-8.
67. Tronchin G, Akudjedu TN, Ahmed M, et al. Progressive subcortical volume loss in treatment-resistant schizophrenia patients after commencing clozapine treatment. *Neuropsychopharmacology* 2020;45:1353-61.
68. Ahmed M, Cannon DM, Scanlon C, et al. Progressive brain atrophy and cortical thinning in schizophrenia after commencing clozapine treatment. *Neuropsychopharmacology* 2015;40:2409-17.
69. Romer AL, Elliott ML, Knodt AR, et al. Pervasively thinner neocortex as a transdiagnostic feature of general psychopathology [Epub ahead of print]. *Am J Psychiatry* 2020 Jun 30;appiajp202019090934. doi: 10.1176/appi.ajp.2020.19090934.
70. Cauda F, Nani A, Liloia D, et al. Finding specificity in structural brain alterations through Bayesian reverse inference. *Hum Brain Mapp* 2020;41:4155-4172.
71. Writing Committee for the Attention-Deficit/Hyperactivity Disorder; Autism Spectrum Disorder; Bipolar Disorder; Major Depressive Disorder; Obsessive-Compulsive Disorder; and Schizophrenia ENIGMA Working Groups; et al. Virtual histology of cortical thickness and shared neurobiology in 6 psychiatric disorders. *JAMA Psychiatry* 2020; doi: 10.1001/jamapsychiatry.2020.2694.
72. Chand GB, Dwyer DB, Erus G, et al. Two distinct neuroanatomical subtypes of schizophrenia revealed using machine learning. *Brain* 2020;143:1027-38.
73. Pan Y, Pu W, Chen X, et al. Morphological profiling of schizophrenia: cluster analysis of MRI-based cortical thickness data. *Schizophr Bull* 2020;46:623-32.
74. Kharitonova M, Martin RE, Gabrieli JDE, et al. Cortical gray-matter thinning is associated with age-related improvements on executive function tasks. *Dev Cogn Neurosci* 2013;6:61-71.
75. Squeglia LM, Jacobus J, Sorg SF, et al. Early adolescent cortical thinning is related to better neuropsychological performance. *J Int Neuropsychol Soc* 2013;19:962-70.
76. Hoekzema E, Barba-Müller E, Pozzobon C, et al. Pregnancy leads to long-lasting changes in human brain structure. *Nature Neuroscience* 2017;20:287-96.
77. Schnack HG, van Haren NEM, Brouwer RM, et al. Changes in thickness and surface area of the human cortex and their relationship with intelligence. *Cereb Cortex* 2015;25:1608-17.
78. Bagot RC, van Hasselt FN, Champagne DL, et al. Maternal care determines rapid effects of stress mediators on synaptic plasticity in adult rat hippocampal dentate gyrus. *Neurobiol Learn Mem* 2009;92:292-300.
79. Amato D, Beasley CL, Hahn MK, et al. Neuroadaptations to antipsychotic drugs: insights from pre-clinical and human post-mortem studies. *Neurosci Biobehav Rev* 2017;76:317-35.
80. Telesford QK, Simpson SL, Burdette JH, et al. The brain as a complex system: using network science as a tool for understanding the brain. *Brain Connect* 2011;1:295-308.
81. Komarov MA, Osipov GV, Burtsev MS. Adaptive functional systems: learning with chaos. *Chaos* 2010;20:045119.
82. Baffy G, Loscalzo J. Complexity and network dynamics in physiological adaptation: an integrated view. *Physiol Behav* 2014;131:49-56.
83. Palaniyappan L. Inefficient neural system stabilization: a theory of spontaneous resolutions and recurrent relapses in psychosis. *J Psychiatry Neurosci* 2019;44:367-383.
84. Post RM, Weiss SRB. Emergent properties of neural systems: how focal molecular neurobiological alterations can affect behavior. *Dev Psychopathol* 1997;9:907-29.
85. Cruse H, Schilling M. Mental states as emergent properties. From walking to consciousness. *Open Mind*; 2015. Available: <https://pub.uni-bielefeld.de/record/2713110> (accessed 2020 Sept. 4).
86. Goldman AW, Burmeister Y, Cesnulevicius K, et al. Bioregulatory systems medicine: An innovative approach to integrating the science of molecular networks, inflammation, and systems biology with the patient's autoregulatory capacity? *Front Psychol* 2015;6:225.
87. Borsboom D, Cramer AOJ, Kalis A. Brain disorders? Not really: Why network structures block reductionism in psychopathology research. *Behav Brain Sci* 2018;24:1-54.
88. Castro N, Siew CSQ. Contributions of modern network science to the cognitive sciences: revisiting research spirals of representation and process. *Proc Math Phys Eng Sci* 2020;476:20190825.
89. Newbold DJ, Laumann TO, Hoyt CR, et al. Plasticity and spontaneous activity pulses in disused human brain circuits. *Neuron* 2020;107:580-589.e6.